

12-19-2013

## Surgery for post-vitrectomy cataract

Diana V. Do

*University of Nebraska Medical Center, [diana.do@unmc.edu](mailto:diana.do@unmc.edu)*

Stephen Gichuhi

*University of Nairobi*

Satyanarayana S. Vedula

*Johns Hopkins University*

Barbara S. Hawkins

*Johns Hopkins University*

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: [https://digitalcommons.unmc.edu/com\\_eye\\_articles](https://digitalcommons.unmc.edu/com_eye_articles)



Part of the [Ophthalmology Commons](#)

---

### Recommended Citation

Do, Diana V.; Gichuhi, Stephen; Vedula, Satyanarayana S.; and Hawkins, Barbara S., "Surgery for post-vitrectomy cataract" (2013). *Journal Articles: Ophthalmology*. 14.

[https://digitalcommons.unmc.edu/com\\_eye\\_articles/14](https://digitalcommons.unmc.edu/com_eye_articles/14)

This Article is brought to you for free and open access by the Ophthalmology at DigitalCommons@UNMC. It has been accepted for inclusion in Journal Articles: Ophthalmology by an authorized administrator of DigitalCommons@UNMC. For more information, please contact [digitalcommons@unmc.edu](mailto:digitalcommons@unmc.edu).

# Surgery for post-vitrectomy cataract (Review)

Do DV, Gichuhi S, Vedula SS, Hawkins BS



**THE COCHRANE  
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2013, Issue 12

<http://www.thecochranelibrary.com>

**WILEY**

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
BACKGROUND . . . . .	2
OBJECTIVES . . . . .	4
METHODS . . . . .	4
RESULTS . . . . .	6
Figure 1. . . . .	7
DISCUSSION . . . . .	8
AUTHORS' CONCLUSIONS . . . . .	8
ACKNOWLEDGEMENTS . . . . .	9
REFERENCES . . . . .	9
CHARACTERISTICS OF STUDIES . . . . .	11
DATA AND ANALYSES . . . . .	13
APPENDICES . . . . .	13
WHAT'S NEW . . . . .	15
HISTORY . . . . .	16
CONTRIBUTIONS OF AUTHORS . . . . .	16
DECLARATIONS OF INTEREST . . . . .	17
SOURCES OF SUPPORT . . . . .	17
INDEX TERMS . . . . .	17

[Intervention Review]

## Surgery for post-vitrectomy cataract

Diana V Do<sup>1</sup>, Stephen Gichuhi<sup>2</sup>, Satyanarayana S Vedula<sup>3</sup>, Barbara S Hawkins<sup>4</sup>

<sup>1</sup>Stanley M. Truhlsen Eye Institute, University of Nebraska Medical Center, Omaha, Nebraska, USA. <sup>2</sup>Department of Ophthalmology, University of Nairobi, Nairobi, Kenya. <sup>3</sup>Johns Hopkins University, Baltimore, Maryland, USA. <sup>4</sup>Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

Contact address: Diana V Do, Stanley M. Truhlsen Eye Institute, University of Nebraska Medical Center, 3902 Leavenworth Street, Omaha, Nebraska, 68105, USA. [diana.do@unmc.edu](mailto:diana.do@unmc.edu).

**Editorial group:** Cochrane Eyes and Vision Group.

**Publication status and date:** New search for studies and content updated (no change to conclusions), published in Issue 12, 2013.

**Review content assessed as up-to-date:** 22 May 2013.

**Citation:** Do DV, Gichuhi S, Vedula SS, Hawkins BS. Surgery for post-vitrectomy cataract. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No.: CD006366. DOI: 10.1002/14651858.CD006366.pub3.

Copyright © 2013 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

### ABSTRACT

#### Background

Cataract formation or acceleration can occur after intraocular surgery, especially following vitrectomy, a surgical technique for removing the vitreous which is used in the treatment of disorders that affect the posterior segment of the eye. The underlying problem that led to vitrectomy may limit the benefit from cataract surgery.

#### Objectives

The objective of this review was to evaluate the effectiveness and safety of surgery for post-vitrectomy cataract with respect to visual acuity, quality of life, and other outcomes.

#### Search methods

We searched CENTRAL (which contains the Cochrane Eyes and Vision Group Trials Register) (*The Cochrane Library* 2013, Issue 4), Ovid MEDLINE, Ovid MEDLINE in-Process and Other Non-Indexed Citations, Ovid MEDLINE Daily Update, Ovid OLDMEDLINE (January 1946 to May 2013), EMBASE (January 1980 to May 2013, Latin American and Caribbean Health Sciences Literature Database (LILACS) (January 1982 to May 2013), PubMed (January 1946 to May 2013), the *meta*Register of Controlled Trials (*mRCT*) ([www.controlled-trials.com](http://www.controlled-trials.com)), ClinicalTrials.gov ([www.clinicaltrial.gov](http://www.clinicaltrial.gov)) and the WHO International Clinical Trials Registry Platform (ICTRP) ([www.who.int/ictrp/search/en](http://www.who.int/ictrp/search/en)). We did not use any date or language restrictions in the electronic searches for trials. We last searched the electronic databases on 22 May 2013.

#### Selection criteria

We planned to include randomized and quasi-randomized controlled trials comparing cataract surgery with no surgery in adult patients who developed cataract following vitrectomy.

#### Data collection and analysis

Two authors screened the search results independently according to the standard methodological procedures expected by The Cochrane Collaboration.

#### Main results

We found no randomized or quasi-randomized controlled trials comparing cataract surgery with no cataract surgery for patients who developed cataracts following vitrectomy surgery.

---

**Surgery for post-vitrectomy cataract (Review)**

Copyright © 2013 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

1

## Authors' conclusions

There is no evidence from randomized or quasi-randomized controlled trials on which to base clinical recommendations for surgery for post-vitrectomy cataract. There is a clear need for randomized controlled trials to address this evidence gap. Such trials should stratify participants by their age, the retinal disorder leading to vitrectomy, and the status of the underlying disease process in the contralateral eye. Outcomes assessed in such trials may include gain of vision on the Early Treatment Diabetic Retinopathy Study (ETDRS) scale, quality of life, and adverse events such as posterior capsular rupture. Both short-term (six-month) and long-term (one-year or two-year) outcomes should be examined.

## PLAIN LANGUAGE SUMMARY

### Surgery to remove cataracts after vitrectomy

#### Review question

We reviewed the evidence about the effect of surgery to remove cataracts (cloudy lens in front of the eye) in people who develop cataracts after vitrectomy, a surgery to remove the vitreous (the clear gel) in the center of the eye.

#### Background

Vitrectomy, a procedure performed for disorders that affect the back part of the eye, can result in the formation or acceleration of cataract. The underlying problem that led to vitrectomy may limit the benefits from cataract surgery.

#### Results

We found no randomized controlled trials (trials in which participants are randomly assigned to one treatment group or another) that evaluated the benefits or risks (or both) of cataract surgery following vitrectomy. Since cataract surgery may lead to loss of vision due to worsening or recurrence of the condition that prompted the vitrectomy, its role in these patients remains uncertain. Future trials to address this review question should separate participants by age, the disorder leading to vitrectomy, and the status of the underlying disease process in the opposite eye. Outcomes relevant to patients such as improvement of vision and quality of life, and harms should be examined both in the short term (six months after surgery) and in the long term (one to two years after surgery).

## BACKGROUND

### Description of the condition

Cataract, an opacification of the crystalline lens in the eye, can be caused by many factors including the natural aging process, metabolic abnormalities, nutritional disorders, chronic ocular inflammation, and trauma. The three types of cataract are classified according to the location of the opacity: cortical, nuclear sclerosis, and posterior subcapsular. Cataract formation or acceleration also can occur after intraocular surgery, especially following vitrectomy, a surgical technique for removing the vitreous that is used in the treatment of disorders that affect the posterior segment of the eye. Vitrectomy causes progression of nuclear sclerotic cataracts. Vitrectomy was first developed by Machemer in 1971 ([Machemer 1971](#)). It is a microsurgical technique in which specialized instru-

ments and techniques are used to gain access to the vitreous cavity and retina. During vitrectomy surgery, three small incisions, each approximately 1.4 mm in length, are made in the eye in order to place instruments: a vitreous cutter, a fiberoptic light source to illuminate the inside of the eye, and an infusion cannula to maintain proper intraocular pressure during the surgery. During the past 40 years, advances in surgical technique and instrumentation have made vitrectomy a common surgical procedure for posterior segment disorders. Vitrectomy is indicated for numerous ocular conditions including vitreous loss in cataract surgery, subluxation of the lens, malignant glaucoma, dense pupillary membranes, non-clearing vitreous hemorrhage due to diabetic retinopathy or vein occlusions, retinal detachment, macular hole, macular pucker, vitreo-macular traction, and endophthalmitis. Although vitrectomy has revolutionized the treatment of posterior segment disorders and improved visual outcomes in patients with

retinal diseases requiring surgical intervention, vitrectomy also is associated with co-morbidities that may compromise visual acuity such as retinal detachment, corneal decompensation, and cataract formation or progression in phakic eyes (Benson 1988). The type of cataract that forms or accelerates after vitrectomy is nuclear sclerotic cataract. Cataract formation or progression is believed to be the most common complication associated with vitrectomy. In fact, in many eyes undergoing vitrectomy, the lens is removed at the same time. Often, the nuclear sclerotic cataracts that develop after vitrectomy limit visual acuity outcomes to a degree that would result in surgical removal of the lens in an otherwise 'normal' eye. The exact pathogenesis of cataract formation or acceleration after vitrectomy is unknown. Older studies have suggested that light toxicity, oxidation of lens proteins, use of intraocular gas, and length of operative time may be causative factors (Cherfan 1991; de Bustros 1988; Ogura 1991). Newer research suggests that vitrectomy surgery increases oxygen tension within the eye; oxygen exposure has been linked with progressive nuclear sclerotic cataract formation (Holekamp 2005; Palmquist 1988).

## Epidemiology

Although cataract progression is common after vitrectomy, only a few prospective studies have evaluated this occurrence. Do and Hawkins performed a review (unpublished) of the pertinent literature in the PubMed database published from 1966 through 2005. A total of 51 studies were found. The majority of published studies on cataract progression after vitrectomy were retrospective analyses. The reported incidence of cataract was highly variable, from 6% to 100% of cases, depending upon the condition that prompted vitrectomy, duration of follow-up, and the method used to monitor development of cataract. These retrospective studies are limited by the non-uniformity of the lens grading system used or the absence of a description of the lens grading system in the published report.

The Vitrectomy for Macular Hole Study, a randomized controlled trial that evaluated vitrectomy for the treatment of macular holes, retrospectively examined the incidence of cataract development among 74 eyes of patients that participated in the study (Cheng 2001). Investigators used a scoring system similar to the Lens Opacities Classification System II, which contains five grading categories for nuclear and posterior subcapsular opacities. Although duration of surgery did not increase the risk for cataract progression, vitrectomy itself was a risk factor for cataract acceleration; 60/74 (81%) eyes in the surgery cohort had nuclear sclerotic cataract progression at six months of follow-up, compared to only 13/74 (18%) fellow eyes in the control group. By two years, 100% of eyes in the surgery cohort had cataract progression, compared to 8% of control eyes. Similarly, Cherfan and colleagues retrospectively reviewed 100 eyes after vitrectomy for idiopathic macular pucker (Cherfan 1991). After an average follow-up of 29 months (range six to 99 months), 80/100 (80%) eyes in the vitrectomy group and

only 24/100 (24%) fellow eyes had developed a visually significant nuclear sclerotic cataract or had undergone cataract extraction. During the late 1990s, the Submacular Surgery Trials (SST) were initiated to evaluate surgical removal of subfoveal choroidal neovascularization (CNV) compared with observation in patients with age-related macular degeneration (AMD) (SST Group N and Group B), ocular histoplasmosis syndrome (OHS) (SST Group H), and idiopathic CNV (SST Group H) (SST 2004a; SST 2004b; SST 2004c). In these three randomized controlled trials, visually significant cataract was defined as either cataract surgery or lens opacity reported by the SST ophthalmologist to be sufficient to reduce visual acuity by two or more lines in a normal eye. Among the AMD participants in the SST Group N study, 80% of eyes assigned to vitrectomy and surgical removal of their subfoveal CNV developed visually significant cataracts at two years of follow-up. Sixty per cent of eyes had undergone cataract surgery by their last follow-up examination two to four years after enrollment. Among the OHS participants in the SST Group H study, 39% of eyes assigned to vitrectomy developed visually significant cataracts, among which 24% underwent cataract removal. The difference between eyes with AMD and eyes with OHS developing post-vitrectomy cataract is likely due to the median age of the patients. Patients under the age of 50 years are less likely to develop post-surgical accelerated nuclear sclerosis (Melberg 1995). Data from the SST provide the largest and most complete follow-up of eyes undergoing vitrectomy that are at high risk for developing visually significant post-surgical nuclear sclerotic cataracts.

## Presentation and diagnosis

Patients who develop post-vitrectomy cataract present with decreased visual acuity despite anatomic or functional success (or both) of the vitrectomy surgery. Individuals who have undergone vitrectomy may have lower levels of baseline (pre-cataract) visual acuity due to the underlying nature of their retinal pathology; therefore patients with post-vitrectomy cataract are more likely to present with poorer vision than individuals with typical senile cataracts. Diagnosis is made with ocular examination using slit-lamp biomicroscopy.

## Description of the intervention

Cataract surgery, typically using phacoemulsification and intraocular lens implantation, is commonly recommended for individuals with visually significant lens opacities. Two features of post-vitrectomy nuclear sclerosis make affected lenses especially challenging for cataract surgeons to remove. The nucleus tends to be harder than in age-related nuclear sclerosis, requiring longer phacoemulsification time during the procedure. Also, the absence of vitreous in the posterior segment allows for more mobility of the posterior capsule, increasing the risk of capsular rupture. Thus, surgery for post-vitrectomy nuclear sclerotic cataract may have a

higher incidence of complications, although evidence from comparative studies is lacking (Ahfat 2003; Biro 2002).

### How the intervention might work

Patients who develop cataract after vitrectomy may undergo cataract extraction; however, visual acuity and other outcomes after cataract surgery may be poor due to the underlying retinal disorder. Most patients who have vitrectomy surgery have serious underlying problems, as indicated by the reasons for vitrectomy. Furthermore, eyes with post-vitrectomy cataract are at risk of complications that affect all eyes that undergo cataract surgery such as endophthalmitis, cystoid macular edema, etc. Thus, vision is often already impaired before cataract surgery and may remain impaired after cataract surgery. Although cataract surgery in a normal eye typically improves vision, the visual prognosis after surgery for post-vitrectomy cataract is uncertain. It likely depends on the success of treatment for the retinal disorder and avoidance of complications during cataract surgery.

### Why it is important to do this review

The incidence of cataract formation after vitrectomy varies widely and has been reported to be between 6% and 100%. The majority of published studies confirm that a high rate of cataract formation occurs, but few data are available on visual acuity outcomes after cataract removal. The retinal problem that led to vitrectomy may progress or recur. However, peer-reviewed data on outcomes after surgery for post-vitrectomy cataract are scarce. Even in situations in which cataract formation is not due to vitrectomy, visual impairment can still exist despite cataract extraction. The Los Angeles Latino Eye Study (Barañano 2007) published visual acuity outcomes after cataract extraction in adult Latinos and reported that 41% of eyes had visual impairment (defined as a best-corrected visual acuity of 20/40 Snellen equivalent or less). Age-related macular degeneration and diabetic retinopathy accounted for approximately 57% of retinal pathology after cataract extraction. In addition, in eyes that have undergone vitrectomy surgery the absence of vitreous in the posterior segment allows for more mobility of the posterior capsule, increasing the risk of capsular rupture. Surgery for post-vitrectomy nuclear sclerotic cataract may have a higher incidence of complications. A systematic review of outcomes from controlled clinical trials would provide information for adequate counseling of patients and for guiding ophthalmologists' recommendations.

## OBJECTIVES

The objective of this review was to evaluate the effectiveness and safety of surgery for post-vitrectomy cataract with respect to visual acuity, quality of life, and other outcomes.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

We planned to include both randomized and quasi-randomized controlled trials in this review. We considered quasi-randomized trials to be trials that had adopted a method of allocation intended to allocate patients in a random fashion but were not strictly random. Examples include allocation by date of birth, social security number, etc. We were to include trials with at least six months' follow-up to allow for reporting of early adverse effects, even though we planned our primary analyses to focus on outcomes at the end of one year of follow-up.

#### Types of participants

We planned to include trials that enrolled adult participants (age 18 years and over) with cataract that had developed after vitrectomy for any indication except for trauma. However, we planned not to exclude trials that included both adult patients who had post-traumatic vitrectomy and patients who had other indications for vitrectomy, except when outcomes were reported separately by indication. We planned to exclude trials that included only trauma cases, because these patients typically are younger and the pathogenesis of cataract formation is different.

#### Types of interventions

We planned to include trials that compared cataract surgery (of any type) with no surgery in such patients.

#### Types of outcome measures

##### Primary outcomes

Visual acuity improvement after cataract surgery of at least three letters on a logMAR chart, one line on the Snellen chart, or equivalent changes on other scales. While we planned to analyze the outcomes at one year, two years, and at longer time points of follow-up as available from included studies, our primary analysis was to focus on one-year follow-up.

## Secondary outcomes

1. Quality of life measured by a validated scale.
2. Contrast sensitivity: improvement of at least one level, regardless of the manner in which it was measured in included trials.
3. Progression of the condition that was the original indication for vitrectomy in patients with diabetic retinopathy (DR) and age-related macular degeneration (AMD) as defined by standard grading scales such as the Diabetic Retinopathy Scale for DR (ETDRS 1991) and the International Scale for AMD (Bird 1995).

## Adverse outcomes

Specific adverse effects of interest included:

- cystoid macular edema;
- intraocular lens-related complications, including dislocation, difficulty in placing the lens;
- capsular opacification;
- retinal detachment (new or recurrent).

We also planned to summarize all other adverse effects reported in included studies.

## Economic outcomes

We planned to tabulate or summarize data on the costs of procedures, consequences of complications, and any cost-effectiveness data reported in included studies in a narrative fashion.

## Search methods for identification of studies

### Electronic searches

In 2013, we revised the searches of electronic databases from the 2011 update. We searched PubMed and the International Clinical Trials Registry Platform, which had not originally been searched. We searched the Cochrane Central Register of Controlled Trials (CENTRAL) (which contains the Cochrane Eyes and Vision Group Trials Register) 2013, Issue 4, part of *The Cochrane Library*. [www.thecochranelibrary.com](http://www.thecochranelibrary.com) (accessed 22 May 2013), Ovid MEDLINE, Ovid MEDLINE In-Process and Other Non-Indexed Citations, Ovid MEDLINE Daily Update, Ovid OLDMEDLINE (January 1946 to August 2013), EMBASE (January 1980 to May 2013), Latin American and Caribbean Health Sciences Literature Database (LILACS) (January 1982 to May 2013), the *metaRegister of Controlled Trials* (*mRCT*) ([www.controlled-trials.com](http://www.controlled-trials.com)), ClinicalTrials.gov ([www.clinicaltrial.gov](http://www.clinicaltrial.gov)), the WHO International Clinical Trials Registry Platform (ICTRP) ([www.who.int/ictrp/search/en](http://www.who.int/ictrp/search/en)) and PubMed (January 1956 to May 2013). We did not use any date

or language restrictions in the electronic searches for trials. We last searched the electronic databases on 22 May 2013.

See: Appendices for details of search strategies for CENTRAL (Appendix 1), MEDLINE (Appendix 2), EMBASE (Appendix 3), LILACS (Appendix 4), *mRCT* (Appendix 5), ClinicalTrials.gov (Appendix 6), ICTRP (Appendix 7) and PubMed (Appendix 8). The UK Clinical Research Network Portfolio Database and the Australian New Zealand Clinical Trials Registry are no longer being searched for this review.

## Searching other resources

We planned to search the reference lists of included studies and the Science Citation Index - Expanded database to identify any additional trials. We did not search any conference proceedings specifically for the purpose of this review.

## Data collection and analysis

### Selection of studies

Two review authors independently screened the titles and abstracts of all records identified in the electronic and manual searches. We labeled each record as 'A - include', 'B - unsure', or 'C - exclude'. Two authors then screened the full reports of records labeled 'B - unsure'; we relabeled each record as 'A - include' or 'C - exclude' based on consensus after review. We listed studies reported in records labeled 'C - exclude' after full-text review in the table of excluded studies with reasons for exclusion. We planned to assess methodological quality for studies labeled as 'A - include'; however, we did not identify any study labeled as 'A - include' by either of the two review authors.

**We found no trials eligible for inclusion in either the original review or the updated review. The methods described below will be applicable to future updates of the review when trials eligible for inclusion have been conducted and reported.**

### Data extraction and management

Two review authors will independently extract data on basic characteristics of each study (including details of study design, characteristics of participants, interventions, and comparators) and the primary and secondary outcomes onto data collection forms developed in collaboration with the Cochrane Eyes and Vision Group. We will resolve discrepancies by discussion. We will contact authors of included studies for missing data. One review author will enter all data into RevMan 5.2 (RevMan 2013) and another author will verify the data.



### Assessment of risk of bias in included studies

Two review authors, working independently, will assess the included studies for sources of systematic bias according to the guidelines in Chapter 8 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011a). We will evaluate studies for the following criteria: method of randomization, allocation concealment (selection bias), masking of outcome assessment, completeness of outcome data and intention-to-treat analysis (attrition bias), and other potential sources of bias including source of trial funding. We will not assess masking of investigators as the interventions to be compared preclude such efforts. Though an artificial lens placed in the eyes of patients in the intervention group may be recognized by the anatomic outcome assessor, visual acuity testing may have been performed by someone not responsible for examining the eye. Also, quality of life data may have been collected by some method that preserves masking of intervention or outcome assessment, or both. We will judge each criterion as either 'low risk of bias', 'high risk of bias', or 'unclear risk of bias'. We will use information in the *Cochrane Handbook for Systematic Reviews of Interventions* to guide our judgment for each criterion. We will contact authors of studies labeled 'unsure' for clarification. Differences between the two review authors will be resolved by discussion.

### Measures of treatment effect

We will calculate a summary risk ratio for dichotomous outcomes (visual acuity improvement, progression of the condition that was the original indication for vitrectomy, and adverse events). We will calculate the mean difference for continuous outcomes (quality of life, cost-effectiveness, and contrast sensitivity).

### Unit of analysis issues

The unit of analysis will be the eye for vision-related outcomes (visual acuity, contrast sensitivity, progression, and adverse events). For quality of life and economic outcomes, the unit of analysis will be the person.

### Dealing with missing data

We will attempt to contact the investigators of included trials for any missing data. If the investigators do not respond within four weeks, we will extract available data from the published report. We will refer to the guidelines in Chapter 16 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011b) for handling missing data.

### Assessment of heterogeneity

We will examine the design and clinical heterogeneity among included studies by carefully analyzing their characteristics. We also

will examine the degree of overlap in the confidence intervals of the studies. If there is poor overlap, this will be taken to indicate the presence of statistical heterogeneity. We will test for statistical heterogeneity formally using the  $\text{Chi}^2$  test. We also will use the  $I^2$  statistic value to determine the proportion of variation due to heterogeneity among studies, and will consider an  $I^2$  value greater than 50% to indicate substantial statistical heterogeneity.

### Assessment of reporting biases

We will examine a funnel plot to identify any evidence of publication bias when a sufficient number of studies are included, i.e., 10 or more.

### Data synthesis

If no significant heterogeneity is detected, either statistically or if there is a small number of trials in the analysis (three or fewer), we will use a fixed-effect model. If the number of trials is greater than three and no heterogeneity has been detected, we will use a random-effects model.

If significant heterogeneity has been detected, we will not combine results to produce a single summary measure. In this case, we will describe the forest plot in the results section of the review.

### Subgroup analysis and investigation of heterogeneity

We will investigate heterogeneity, if present, through subgroup analyses. If sufficient data are available, we will conduct subgroup analyses based on the agents used to fill the vitreous space after vitrectomy, e.g., air, different gases, and by different indications for vitrectomy.

### Sensitivity analysis

We will conduct sensitivity analyses to determine the impact of exclusion of studies of lower methodological quality, including quasi-randomized trials, and exclusion of industry-funded studies and unpublished studies.

## RESULTS

### Description of studies


#### Results of the search

The original electronic searches retrieved a total of 1949 references and 29 additional records from clinical trials registers. After

independent review of the titles and abstracts by two review authors, we retrieved 36 full-text articles. We found no randomized or quasi-randomized trials eligible for inclusion in the review. An updated search was done in April 2011; we retrieved 785 references and 18 additional records from clinical trials registers. We assessed the records but none were eligible for inclusion in the review. The most recent updated search conducted in May 2013 yielded 1009 references and 120 additional records from clinical trials

registers. We assessed the records but none were eligible for inclusion in the review (Figure 1). During the process of the most recent update, we also re-assessed the eligibility for all previously excluded studies, and removed 22 out of 36 references from the list by reviewing titles and abstracts. All of the 22 references were either non-human study or non randomized controlled trials. We still kept the reasons for exclusion for the remaining 14 references for 10 studies in the 'Characteristics of excluded studies' table.

Figure 1. Study flow diagram.



### Included studies

We did not identify any studies eligible for inclusion in this review.

### Excluded studies

Review of the full-text articles did not identify any studies that are relevant to the objective of this systematic review.

### Risk of bias in included studies

We found no trials eligible for inclusion in the review for assessment of risk of bias.

### Effects of interventions

We found no information on effects of interventions from trials eligible for inclusion in the review.

## DISCUSSION

The majority of the published literature on this subject is limited to retrospective case reports or non-randomized prospective case series (Ahfar 2003). Any attempt to draw conclusions from these non-randomized studies would be misleading. In addition, there is no reliable method to identify all observational studies on this topic; therefore data collection is likely to be incomplete. Thus, our protocol specifically stated that we were interested in outcomes based on randomized or quasi-randomized clinical trials, not observational studies. This dearth of information on surgery for post-vitrectomy cataract indicates that information on this topic is needed, as thousands of patients undergo vitrectomy each year and are at risk of development of cataract and cataract surgery. Documentation of both the risks and benefits of surgery for post-vitrectomy cataract is needed to inform patient counseling and clinical recommendations.

## AUTHORS' CONCLUSIONS

### Implications for practice

In the absence of data from controlled trials, ophthalmologists have no reliable evidence to use when counseling patients regarding the risks and benefits of surgery for post-vitrectomy cataract, and no basis for recommendations for or against cataract surgery or when to intervene surgically. It is possible that some of the complications

of surgery for post-vitrectomy cataract could be reduced if surgery were known to be beneficial and to be beneficial if performed at an early stage of development of nuclear sclerosis. Data from retrospective studies are inadequate for these purposes.

### Implications for research

There is a clear need for well-designed randomized controlled trials to evaluate the benefits and risks of surgery for cataracts that develop following vitrectomy. We recommend that randomization in such trials be stratified by patients' age, retinal disorder leading to vitrectomy, and status of the pathologic process in the contralateral eye. Because patients who undergo vitrectomy already have reduced vision due to the underlying condition that prompted vitrectomy, relevant outcomes such as quality of life should be considered in addition to visual acuity and other clinical measures of vision. We recommend that restoration or gain of eight or more letters vision on the ETDRS scale with cataract surgery would be a reasonable expectation in this patient population. Analyses should include both short-term (six months) and long-term (one-year to two-year) outcomes. Data on adverse outcomes, including complications of surgery such as posterior capsular rupture, should be documented in future trials.

However, there are ethical difficulties in conducting a randomized trial of surgery for post-vitrectomy cataract. Observation of progression of post-vitrectomy cataract may not be considered an ethical alternative. In certain patient populations, such as those with significant central vision loss due to macular scars, observation of post-vitrectomy cataract may be ethical, and conducting a randomized controlled trial to determine if cataract surgery improves quality of life may be a reasonable option. One possible trial design would be to enroll patients over the age of 50 years who are scheduled for vitrectomy, who do not have visually significant cataract in the study eye, and who have visual acuity better than 20/40 in the fellow eye in a two-stage RCT. The initial randomization would be between vitrectomy alone and vitrectomy plus lens removal. Any cataract surgery in the fellow eye would be deferred for at least six months. At six months, study eyes in the deferred group in which visually significant cataract had developed would then be randomized to cataract surgery or deferment for another six months. At the end of the second six-month period, all patients with visually significant cataract in the study eye could be offered surgery. This design would permit evaluation of:

- time to development or progression of cataract in the initial vitrectomy only arm;
- time to progression or recurrence of the underlying retinal problem that led to vitrectomy in the two initial treatment arms;

- comparison of quality of life between the two arms of the original randomization; and
- (with the second randomization) comparison of the above outcomes with and without cataract surgery.

This design would take account of the large number of patients who have lens surgery at the time of vitrectomy and might support that practice or suggest that it was desirable only in selected subgroups of patients, as long as the fellow eye had useful vision.

Concerns about surgical complications, prognosis for recovery of visual acuity, and uncertainty regarding progression of underlying retinal disorder, are important considerations in establishing equipoise necessary for randomization. Further, insufficient infor-

mation on outcomes important to patients, such as vision-related quality of life, indicates an inability to reasonably assess risks and benefits of surgery for post-vitrectomy cataract.

## ACKNOWLEDGEMENTS

We thank Iris Gordon, Trials Search Co-ordinator for the Cochrane Eyes and Vision Group, for designing and conducting the electronic searches. We thank Marie Diener-West for her comments on the protocol and Nancy Holekamp and Roberta Scherer for their comments on the review. The CEVG methodologist Xue Wang (XW) and the Editorial team provided co-ordination and support.

## REFERENCES

### References to studies excluded from this review

#### Ahmadieh 2005 {published data only}

Ahmadieh H, Moradian S, Faghihi H, Parvaresh MM, Ghanbari H, Mehryar M, et al. Anatomic and visual outcomes of scleral buckling versus primary vitrectomy in pseudophakic and aphakic retinal detachment: six-month follow-up results of a single operation--report no. 1. *Ophthalmology* 2005;**112**(8):1421-9.

#### Asaria 2001 {published data only}

Asaria RH, Kon CH, Bunce C, Charteris DG, Wong D, Khaw PT, et al. Adjuvant 5-fluorouracil and heparin prevents proliferative vitreoretinopathy: results from a randomized, double-blind, controlled clinical trial. *Ophthalmology* 2001;**108**(7):1179-83.

#### Batman 1999 {published data only}

Batman C, Cekic O. Vitrectomy with silicone oil or long-acting gas in eyes with giant retinal tears: long-term follow-up of a randomized clinical trial. *Retina* 1999;**19**(3):188-92.

#### Blankenship 1980 {published data only}

Blankenship GW. The lens influence on diabetic vitrectomy results. Report of a prospective randomized study. *Archives of Ophthalmology* 1980;**98**(12):2196-8.

#### DRVS 1985 {published data only}

The Diabetic Retinopathy Vitrectomy Study Research Group. Early vitrectomy for severe proliferative diabetic retinopathy in eyes with useful vision. Results of a randomized trial--Diabetic Retinopathy Vitrectomy Study Report 3. The Diabetic Retinopathy Vitrectomy Study Research Group. *Ophthalmology* 1988;**95**(10):1307-20. The Diabetic Retinopathy Vitrectomy Study Research Group. Early vitrectomy for severe vitreous hemorrhage in diabetic retinopathy. Four-year results of a randomized trial:

Diabetic Retinopathy Vitrectomy Study Report 5. *Archives of Ophthalmology* 1990;**108**(7):958-64.

The Diabetic Retinopathy Vitrectomy Study Research Group. Early vitrectomy for severe vitreous hemorrhage in diabetic retinopathy. Two-year results of a randomized trial. Diabetic Retinopathy Vitrectomy Study report 2. The Diabetic Retinopathy Vitrectomy Study Research Group. *Archives of Ophthalmology* 1985;**103**(11):1644-52.

#### Ehud 1988 {published data only}

Ehud A, Varda C, Joseph M, Giora T. Management of complicated retinal detachment by vitrectomy and silicone oil injection. *Metabolic, Pediatric, and Systemic Ophthalmology* 1988;**11**(1-2):63-6.

#### Ezra 2004 {published data only}

Ezra E, Gregor ZJ. Surgery for idiopathic full-thickness macular hole: two-year results of a randomized clinical trial comparing natural history, vitrectomy, and vitrectomy plus autologous serum: Morfields Macular Hole Study Group RAeport no. 1. *Archives of Ophthalmology* 2004;**122**(2):224-36.

#### Freeman 1997 {published data only}

Freeman WR, Azen SP, Kim JW, el-Haig W, Mishell DR 3rd, Bailey I. Vitrectomy for the treatment of full-thickness stage 3 or 4 macular holes. Results of a multicentered randomized clinical trial. The Vitrectomy for Treatment of Macular Hole Study Group. *Archives of Ophthalmology* 1997;**115**(1):11-21.

#### Haimann 1984 {published data only}

Haimann MH, Abrams GW. Prevention of lens opacification during diabetic vitrectomy. *Ophthalmology* 1984;**91**(2):116-21.

#### Silicone Study 1992 {published data only}

Abrams GW, Azen SP, McCuen BW 2nd, Flynn HW Jr, Lai MY, Ryan SJ. Vitrectomy with silicone oil or long-acting gas in eyes with severe proliferative vitreoretinopathy: results of

additional and long-term follow-up. Silicone Study report 11. *Archives of Ophthalmology* 1997;**115**(3):335–44.

Anonymous. Vitrectomy with silicone oil or perfluoropropane gas in eyes with severe proliferative vitreoretinopathy: results of a randomized clinical trial. Silicone Study Report 2. *Archives of Ophthalmology* 1992; **110**(6):780–92.

Anonymous. Vitrectomy with silicone oil or sulfur hexafluoride gas in eyes with severe proliferative vitreoretinopathy: results of a randomized clinical trial. Silicone Study Report 1. *Archives of Ophthalmology* 1992; **110**(6):770–9.

## Additional references

### Ahfat 2003

Ahfat FG, Yuen CH, Groenewald CP. Phacoemulsification and intraocular lens implantation following pars plana vitrectomy: a prospective study. *Eye* 2003;**17**(1):16–20.

### Barañano 2007

Barañano AE, Wu J, Mazhar K, Azen SP, Varma R, Los Angeles Latino Eye Study Group. Visual acuity outcomes after cataract extraction in adult Latinos. The Los Angeles Latino Eye Study. *Ophthalmology* 2008;**115**(5):815–21.

### Benson 1988

Benson WE. Vitrectomy in clinical ophthalmology. In: Duane TD editor(s). *Duane's Clinical Ophthalmology*. Vol. 5, Philadelphia: Lippincott, 1988:15–7.

### Bird 1995

Bird AC, Bressler NM, Bressler SB, Chisholm IH, Coscas G, Davis MD, et al. An international classification and grading system for age-related maculopathy and age-related macular degeneration. The International ARM Epidemiological Study Group. *Survey of Ophthalmology* 1995;**39**(5):367–74.

### Biro 2002

Biro Z, Kovacs B. Results of cataract surgery in previously vitrectomized eyes. *Journal of Cataract and Refractive Surgery* 2002;**28**(6):1003–6.

### Cheng 2001

Cheng L, Azen SP, El-Bradey MH, Scholz BM, Chaidhawangul S, Toyoguchi M, et al. Duration of vitrectomy and postoperative cataract in the vitrectomy for macular hole study. *American Journal of Ophthalmology* 2001;**132**(6):881–7.

### Cherfan 1991

Cherfan GM, Michels RG, de Bustros S, Enger C, Glaser BM. Nuclear sclerotic cataract after vitrectomy for idiopathic epiretinal membranes causing macular pucker. *American Journal of Ophthalmology* 1991;**111**(4):434–8.

### de Bustros 1988

de Bustros S, Thompson JT, Michels RG, Enger C, Rice TA, Glaser BM. Nuclear sclerosis after vitrectomy for idiopathic epiretinal membrane. *American Journal of Ophthalmology* 1988;**105**(2):160–4.

### ETDRS 1991

Anonymous. Fundus photographic risk factors for progression of diabetic retinopathy. ETDRS report number 12. Early Treatment Diabetic Retinopathy Study Research Group. *Ophthalmology* 1991;**98**(Suppl 5):823–33.

### Higgins 2011a

Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

### Higgins 2011b

Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: Special topics in statistics. In: Higgins JPT, Green S (editors), *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

### Holekamp 2005

Holekamp NM, Shui YB, Beebe DC. Vitrectomy surgery increases oxygen exposure to the lens: a possible mechanism for nuclear cataract formation. *American Journal of Ophthalmology* 2005;**139**(2):302–10.

### Machemer 1971

Machemer R, Buettner H, Norton EWD, Parel JM. Vitrectomy: A pars plana approach. *Transactions of the American Academy of Ophthalmology and Otolaryngology* 1971;**75**(4):813–20.

### Melberg 1995

Melberg NS, Thomas MA. Nuclear sclerotic cataract after vitrectomy in patients under 50 years of age. *Ophthalmology* 1995;**102**(10):1466–71.

### Ogura 1991

Ogura Y, Takanashi T, Ishigooka H, Ogino N. Quantitative analysis of lens changes after vitrectomy by fluorophotometry. *American Journal of Ophthalmology* 1991;**111**(2):179–83.

### Palmquist 1988

Palmquist BM, Philipson B, Fagerholm P. Nuclear cataract - a microradiographic study. *Acta Ophthalmologica* 1988;**66**(6):671–7.

### RevMan 2013

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). 5.2. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2013.

### SST 2004a

Hawkins BS, Bressler NM, Miskala PH, Bressler SB, Holekamp NM, Marsh MJ, et al. Surgery for subfoveal choroidal neovascularization in age-related macular degeneration: ophthalmic findings: SST report no.11. *Ophthalmology* 2004;**111**(11):1967–80.

### SST 2004b

Bressler NM, Bressler SB, Childs AL, Haller JA, Hawkins BS, Lewis H, et al. Surgery for hemorrhagic choroidal

neovascular lesions of age-related macular degeneration: Ophthalmic findings. SST report no. 13. *Ophthalmology* 2004;**111**(11):1993–2006.

**SST 2004c**

Hawkins BS, Bressler NM, Bressler SB, Davidorf FH, Hoskins JC, Marsh MJ, et al. Surgical removal vs observation for subfoveal choroidal neovascularization, either associated with the ocular histoplasmosis syndrome or idiopathic: I. Ophthalmic findings from a randomized clinical trial: Submacular Surgery Trials (SST) Group H Trial: SST Report No. 9. *Archives of Ophthalmology* 2004;**122**(11): 1597–611.

**References to other published versions of this review**

**Do 2007**

Do DV, Hawkins BS, Gichuhi S. Surgery for post-vitrectomy cataract. *Cochrane Database of Systematic Reviews* 1, Issue 2007. [DOI: 10.1002/14651858.CD006366]

**Do 2008**

Do DV, Hawkins B, Gichuhi S, Vedula SS. Surgery for post-vitrectomy cataract. *Cochrane Database of Systematic Reviews* 2008, Issue 3. [DOI: 10.1002/14651858.CD006366.pub2]

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Ahmadieh 2005	RCT, not post-vitrectomy patients
Asaria 2001	RCT, post-vitrectomy patients, not for cataract surgery
Batman 1999	RCT, not post-vitrectomy patients, not cataract surgery
Blankenship 1980	RCT, not post-vitrectomy patients, not for cataract surgery
DRVS 1985	RCT or CCT, not post-vitrectomy patients, not cataract surgery
Ehud 1988	RCT, not post-vitrectomy patients, not for cataract surgery
Ezra 2004	RCT, not post-vitrectomy patients, not for cataract surgery
Freeman 1997	RCT, not post-vitrectomy patients, not for cataract surgery
Haimann 1984	RCT, not post-vitrectomy patients, not for cataract surgery
Silicone Study 1992	RCT for post-vitrectomy patients, not for cataract surgery

CCT: controlled clinical trial

RCT: randomized controlled trial

## DATA AND ANALYSES

This review has no analyses.

## APPENDICES

### Appendix 1. CENTRAL search strategy

- #1 MeSH descriptor: [Vitreotomy] explode all trees
- #2 MeSH descriptor: [Vitreoretinal Surgery] explode all trees
- #3 (vitrectom\* or post-vitrectom\* or postvitrectom\*)
- #4 (vitreous surg\* or vitreous resection\* or vitreoretinal surg\*)
- #5 #1 or #3 or #4
- #6 MeSH descriptor: [Cataract] explode all trees
- #7 MeSH descriptor: [Cataract Extraction] explode all trees
- #8 MeSH descriptor: [Capsulorhexis] explode all trees
- #9 MeSH descriptor: [Phacoemulsification] explode all trees
- #10 cataract\*
- #11 lens\*
- #12 (phakectom\* or zonulolys\* or cataractom\*)
- #13 (pha?oemulsif\* or pha?o or Capsulor?hexis)
- #14 #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13
- #15 #5 and #14

### Appendix 2. MEDLINE (OvidSP) search strategy

1. Randomized Controlled Trial.pt.
2. Controlled Clinical Trial.pt.
3. (randomized or randomised).ab,ti.
4. placebo.ab,ti.
5. drug therapy.fs.
6. randomly.ab,ti.
7. trial.ab,ti.
8. groups.ab,ti.
9. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8
10. exp animals/ not humans.sh.
11. 9 not 10
12. exp Vitrectomy/
13. exp Vitreoretinal Surgery/
14. (vitrectom\* or post-vitrectom\* or postvitrectom\*).tw.
15. (vitreous surg\* or vitreous resection\* or vitreoretinal surg\*).tw.
16. 12 or 14
17. exp Cataract/
18. exp Cataract Extraction/
19. exp Capsulorhexis/
20. exp Phacoemulsification/
21. cataract\*.tw.
22. lens\*.tw.
23. (Phakectom\* or Zonulolys\* or cataractom\*).tw.



- 24. (pha?oemulsif\* or pha#o or Capsulor?hexis).tw.
- 25. or/17-24
- 26. 16 and 25
- 27. 11 and 26

### Appendix 3. EMBASE.com search strategy

- #1 'randomized controlled trial'/exp
- #2 'randomization'/exp
- #3 'double blind procedure'/exp
- #4 'single blind procedure'/exp
- #5 random\*:ab,ti
- #6 #1 OR #2 OR #3 OR #4 OR #5
- #7 'animal'/exp OR 'animal experiment'/exp
- #8 'human'/exp
- #9 #7 AND #8
- #10 #7 NOT #9
- #11 #6 NOT #10
- #12 'clinical trial'/exp
- #13 (clin\* NEAR/3 trial\*):ab,ti
- #14 ((singl\* OR doubl\* OR trebl\* OR tripl\*) NEAR/3 (blind\* OR mask\*)):ab,ti
- #15 'placebo'/exp
- #16 placebo\*:ab,ti
- #17 random\*:ab,ti
- #18 'experimental design'/exp
- #19 'crossover procedure'/exp
- #20 'control group'/exp
- #21 'latin square design'/exp
- #22 #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21
- #23 #22 NOT #10
- #24 #23 NOT #11
- #25 'comparative study'/exp
- #26 'evaluation'/exp
- #27 'prospective study'/exp
- #28 control\*:ab,ti OR prospectiv\*:ab,ti OR volunteer\*:ab,ti
- #29 #25 OR #26 OR #27 OR #28
- #30 #29 NOT #10
- #31 #30 NOT (#11 OR #23)
- #32 #11 OR #24 OR #31
- #33 'vitrectomy'/exp
- #34 'vitreoretinal surgery'/de
- #35 vitrectom\*:ab,ti OR postvitrectom\*:ab,ti
- #36 (vitreous NEAR/1 surg\*):ab,ti OR (vitreous NEAR/1 resection\*):ab,ti OR (vitreoretinal NEAR/1 surg\*):ab,ti
- #37 #33 OR #34 OR #35 OR #36
- #38 'cataract'/exp
- #39 'cataract extraction'/exp
- #40 'capsulorhexis'/exp
- #41 'phacoemulsification'/exp
- #42 cataract\*:ab,ti
- #43 lens\*:ab,ti
- #44 phakectom\*:ab,ti OR zonulolys\*:ab,ti OR cataractom\*:ab,ti
- #45 pha?oemulsif\* OR pha?o OR capsulor?hexis

#46 #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45  
#47 #32 AND #37 AND #46

#### **Appendix 4. LILACS search strategy**

(Vitrecom\$ OR post-vitrecom\$ OR postvitrectom\$ OR “vitreous surgery” OR “vitreous surgeries” OR “vitreous resection” OR “vitreoretinal surgery” OR “vitreoretinal surgeries” OR MH:E04.540.960\$) AND (Cataract\$ OR Catarata\$ OR MH:C11.510.245\$ OR MH:E04.540.825.249\$ OR MH:E04.943.875\$ OR lens\$ OR Capsulorhexis OR Capsulorrexix OR Capsulorrexix OR Capsulorrhexis OR Phacoemulsif\$ OR Facoemulsif\$ OR Phakectom\$ OR Zonulolys\$ OR catarectom\$ OR Phakoemulsif\$)

#### **Appendix 5. metaRegister of Controlled Trials search strategy**

cataract and vitrectomy

#### **Appendix 6. ClinicalTrials.gov search strategy**

cataract AND vitrectomy

#### **Appendix 7. ICTRP search strategy**

cataract AND vitrectomy

#### **Appendix 8. PubMed search strategy**

#1 ((randomized controlled trial[pt]) OR (controlled clinical trial[pt]) OR (randomised[tiab] OR randomized[tiab]) OR (placebo[tiab]) OR (drug therapy[sh]) OR (randomly[tiab]) OR (trial[tiab]) OR (groups[tiab])) NOT (animals[mh] NOT humans[mh])  
#2 (vitrectom\*[tiab] OR post-vitrecom\*[tiab] OR postvitrectom\*[tiab]) NOT Medline[sb]  
#3 (“vitreous surgery”[tiab] OR “vitreous surgeries”[tiab] OR vitreous resection\*[tiab] OR “vitreoretinal surgery”[tiab] OR “vitreoretinal surgery”[tiab]) NOT Medline[sb]  
#4 (#2 OR #3)  
#5 (cataract\*[tiab]) NOT Medline[sb]  
#6 (lens\*[tiab]) NOT Medline[sb]  
#7 (Phakectom\*[tiab] OR Zonulolys\*[tiab] OR catarectom\*[tiab]) NOT Medline[sb]  
#8 (phaco\*[tiab] OR phako\*[tiab] OR Capsulorhexis[tiab] OR Capsulorrhexis[tiab]) NOT Medline[sb]  
#9 (#5 OR #6 OR #7 OR #8)  
#8 (#1 AND #4 AND #9)

## **WHAT'S NEW**

Last assessed as up-to-date: 22 May 2013.

Date	Event	Description
2 December 2013	New citation required but conclusions have not changed	Issue 12, 2013: Some sections of the review have been updated including the 'Implications for practice' section.
2 December 2013	New search has been performed	Issue 12, 2013: Updated searches yielded no new trials.

## HISTORY

Protocol first published: Issue 1, 2007

Review first published: Issue 3, 2008

Date	Event	Description
5 May 2011	New search has been performed	Issue 6, 2011: Updated searches yielded no new trials.
26 March 2008	Amended	Converted to new review format.
27 September 2006	New citation required and conclusions have changed	Substantive amendment.

## CONTRIBUTIONS OF AUTHORS

Conceiving the review: BSH

Designing the review: DVD, BSH

Co-ordinating the review: BSH

Data collection for the review

- Designing electronic search strategies: CEVG Trials Search Co-ordinator, DVD, BSH, SG, SV

- Undertaking searches: CEVG Trials Search Co-ordinator, DVD, BSH, SG, SV

- Screening search results: DVD, BSH, SG, SV

- Organizing retrieval of papers: DVD, XW

- Screening retrieved papers against inclusion criteria: DVD, BSH, SV, XW

- Appraising quality of papers: DVD, BSH

- Extracting data from papers: DVD, BSH, SV

- Writing to authors of papers for additional information: DVD, BSH

- Providing additional data about papers: BSH, SG

- Data management for the review: SG, SV

- Entering data into RevMan: SG, DVD, SV

**Surgery for post-vitrectomy cataract (Review)**

Copyright © 2013 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Analysis of data: BSH, DVD, SV

Interpretation of data

- Providing a methodological perspective: BSH, SG, SV

- Providing a clinical perspective: DVD, BSH, SG, SV

- Providing a policy perspective: BSH, DVD, SG, SV

Writing the review: DVD, BSH, SG, SV

Securing funding for the review: BSH, DVD

Performing previous work that was the foundation of the current study: BSH, DVD

Updating the review: BSH, DVD, XW

Guarantor of the review: DVD

## DECLARATIONS OF INTEREST

None known.

## SOURCES OF SUPPORT

### Internal sources

- Research to Prevent Blindness Grant to Wilmer Eye Institute, USA.
- Retina Research Fund, Wilmer Eye Institute, The Johns Hopkins Hospital, Baltimore, USA.

### External sources

- National Eye Institute, National Institutes of Health, USA.

Methodological support from Cochrane Eyes and Vision - US Project is funded by Grant 1 U01 EY020522-01

## INDEX TERMS

### Medical Subject Headings (MeSH)

\*Cataract Extraction; Cataract [\*etiology]; Quality of Life; Randomized Controlled Trials as Topic; Visual Acuity; Vitrectomy [\*adverse effects]

### MeSH check words

Adult; Humans