

2-17-2018

## Effect of Nitrosamine (NNAT) on Embryogenesis: Evidence from a Study Using Avian Embryos Exposed to NNAT

Moses New-Aaron

*University of Nebraska Medical Center, [moses.newaaron@unmc.edu](mailto:moses.newaaron@unmc.edu)*

Martha Rhoades

*University of Nebraska - Lincoln*

Jane L. Meza

*University of Nebraska Medical Center, [jmeza@unmc.edu](mailto:jmeza@unmc.edu)*

Jeffrey Wallman

*University of Nebraska - Lincoln*

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: [https://digitalcommons.unmc.edu/coph\\_pres](https://digitalcommons.unmc.edu/coph_pres)



Part of the [Public Health Commons](#)

---

### Recommended Citation

New-Aaron, Moses; Rhoades, Martha; Meza, Jane L.; and Wallman, Jeffrey, "Effect of Nitrosamine (NNAT) on Embryogenesis: Evidence from a Study Using Avian Embryos Exposed to NNAT" (2018). *Posters and Presentations: College of Public Health*. 9.

[https://digitalcommons.unmc.edu/coph\\_pres/9](https://digitalcommons.unmc.edu/coph_pres/9)

This Poster is brought to you for free and open access by the College of Public Health at DigitalCommons@UNMC. It has been accepted for inclusion in Posters and Presentations: College of Public Health by an authorized administrator of DigitalCommons@UNMC. For more information, please contact [digitalcommons@unmc.edu](mailto:digitalcommons@unmc.edu).

Moses New-Aaron MPH,BMLS<sup>1</sup>, Martha Rhoades PhD<sup>2</sup>, Jeffery Wallman BS<sup>2</sup>, Chris Wichman PhD<sup>1</sup>, Jane Meza PhD<sup>1</sup>

1. Department of Biostatistics, University of Nebraska Medical Center, Omaha,

2. School of Natural Resources, University of Nebraska-Lincoln

Research was conducted at School of Natural Resources, UNL and the Department of Biostatistics, UNMC

## Background

- *N*-nitrosoatrazine (NNAT) forms in the acidic environment of the human stomach when nitrite (from nitrate) and atrazine are present together<sup>1</sup>.
- NNAT is a nitrosamine, many of which are known carcinogens, but little is known about NNAT toxicity. The effects of NNAT on embryo development are virtually unknown<sup>2</sup>.
- Chromosomal aberrations and an increase in mitotic index were observed in human lymphocytes exposed to NNAT at 1,000 to 10,000-fold lower concentrations than nitrate, nitrite, or atrazine alone<sup>3</sup> and NNAT has been shown to be mutagenic<sup>4</sup>.
- We were among the first to account for the correlation of exposure to agricultural mixtures from drinking water to birth defect rates in the 93 counties of Nebraska<sup>5</sup>

## Hypothesis

We hypothesize that the embryos exposed to NNAT would have delayed development and increased mortality compared to unexposed embryos.

## Method

### Treatment, Incubation, and Analysis of Chicken Embryos

➤ For dosing, the eggshell was punctured on the broad, flat side of the egg. Eggs were administered 50  $\mu$ L doses of NNAT in Dimethyl Sulfoxide (DMSO). Untreated eggs served as negative controls and eggs injected with DMSO served as vehicle controls. DMSO was controlled for by eggs injected with distilled water

➤ The different doses of NNAT administered were 0.245, 1.11, 2.22 and 3.33

➤ 330 eggs were incubated for 5 days in a forced air incubator at 38 °C and 65-75% relative humidity

### Data collection and analysis

➤ After 5 days of incubating the treated eggs the weight, morphology, and vital status of the embryos were obtained

➤ All data analysis was done using SAS 9.4

➤ It was observed that the position of the eggs in the incubator affected their weight from a linear regression analysis

➤ It was also observed from a Chi square analysis that the column arrangement of the eggs was associated to the vital status of the eggs.

➤ The effects of NNAT treatment on the weight of the embryos was obtained after controlling the effect of position in a centered ANCOVA

➤ The likelihood of death after treatment with NNAT was obtained after adjusting for column of the eggs in a multiple logistic regression

➤ The lethal dose of NNAT was determined using PROC PROBIT with a log<sub>10</sub> transformation using SAS 9.4

## Findings

Fig1: Weight of treated embryos




Fig. 2: Vital Status of the treated embryos




Table 1: Effect of treatment on the weight of embryo

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	9	13.45744989	1.49527221	37.21	<.0001
Error	241	9.68413811	0.04018315		
Uncorrected Total	250	23.14158800			

Table 2: Mean weight of the treated embryos

Parameter	Estimate	Standard Error	t Value	Pr >  t
Blank	0.3430998831	0.02255601	15.21	<.0001
Water	0.1764897659	0.02955875	5.97	<.0001
50:50 Water/DMSO	0.1815501283	0.06339472	2.86	0.0046
DMSO	0.1261959807	0.02655123	4.75	<.0001
0.245NNAT in DMSO	0.1566642152	0.06339753	2.47	0.0142
1.11NNAT in DMSO	0.1280735693	0.04603997	2.78	0.0058
2.22NNAT in DMSO	0.1083519218	0.04603250	2.35	0.0194
3.33NNAT in DMSO	0.1001346495	0.06342173	1.58	0.1157
center	0.0045704343	0.00103479	4.42	<.0001


Table 3: Comparisons of the weight of the embryos among the treatment groups

Treat_A_Incr Comparison	Difference Between Means	Simultaneous 95% Confidence Limits
0.245NNAT/DMSO - 1.11NNAT/DMSO	0.03398	-0.20549 0.27345
0.245NNAT/DMSO - 2.22NNAT/DMSO	0.05298	-0.18649 0.29245
0.245NNAT/DMSO - 3.33NNAT/DMSO	0.06110	-0.21302 0.33522
DMSO - Blank	-0.21529	-0.32181 -0.10876 ***
DMSO - Water	-0.05208	-0.17357 0.06941
1.11NNAT/DMSO - Blank	-0.22311	-0.37973 -0.06649 ***
2.22NNAT/DMSO - Blank	-0.24211	-0.39873 -0.08549 ***
3.33NNAT/DMSO - Blank	-0.25023	-0.45597 -0.04450 ***

Table 4: Likelihood of embryo death after treatment with NNAT

Effect	Odds Ratio Estimates		
	Point Estimate	95% Wald Confidence Limits	
Blank vs DMSO	0.351	0.136	0.911
Water vs DMSO	0.249	0.068	0.909
1.11NNAT in DMSO vs DMSO	1.416	0.505	3.970
2.22NNAT in DMSO vs DMSO	2.905	1.167	7.228
3.33NNAT in DMSO vs DMSO	3.344	1.320	8.475
column	1.074	0.924	1.249

Fig. 3: Predicted probabilities for lethal dose of NNAT



## Discussion

- It was found that the weight of the embryos differ significantly in at least 2 of the treatment groups ( $p < 0.0001$ ) (Table 1)
- NNAT decreased the weight of the embryos (0.245NNAT Wgt=0.16g,  $p=0.014$ , 1.11NNAT Wgt=0.13g,  $p=0.0058$ , 2.22NNAT Wgt=0.11g,  $p=0.019$ ) ( Fig1, Table 2)
- Embryos treated with DMSO had lower weight compared to the embryos treated with water and the untreated embryos (Fig1)
- The mean weight of embryos treated with NNAT differ significantly when compared to the untreated embryo ( $P < 0.05$ ) (Table 3)
- Higher mortality for embryos exposed to NNAT compared to unexposed embryos. (Fig 2)
- NNAT-exposed embryos were more likely to die compared to unexposed embryo (2.22NNAT vs. DMSO(OR: 2.91, 95% C.I. 1.17,7.23) and 3.33NNAT vs. DMSO (OR=3.34, 95% C.I. 1.32,8.47) (Table 4)
- The dose of NNAT required to kill 50% of the embryo is  $3.07 \mu\text{mol/l}$  (95 C.I. 2.16, 7.14) (Fig 3)

## Conclusion and Future Directions

- Developing chicken embryos were used in the present study to evaluate the teratogenic potential of *N*-nitrosoatrazine, a nitrosated form of the agricultural atrazine
- NNAT has some teratogenic properties based on the weight and survival of chicken embryos
- A limitation to achieving these objectives is the low solubility of NNAT in water (290 mg/L) and the unclear toxic nature of DMSO to living tissues.
- A more suitable solvent for demonstrating the teratogenic properties of NNAT should be considered in future studies

## References

1. Garcia A, Morales P, Arranz N, Delgado E, Rafter J, & Haza AI. (2008). Induction of apoptosis and reactive oxygen species production by *N*-nitrosopiperidine and *N*-nitrosodibutylamine in human leukemia cells. *Journal of Applied Toxicology*, 28, 455-465.
2. Hamburger V & Hamilton HL. (1992). A Series of Normal Stages in the Development of the Chick Embryo. *Developmental Dynamics*, 195, 231-272.
3. Hanna-Attisha M, LaChance J, Sadler RC, & Schnepf AC. (2016). Elevated blood lead levels in children associated with the Flint drinking water crisis: a spatial analysis of risk and public health response. *American Journal of Public Health*, 106, 283-290.
4. Joo H, Choi K, & Hodgson E. (2010). Human metabolism of atrazine. *Pesticide Biochemistry and Physiology*, 98, 73-79.
5. New-Aaron, M., Rhoades, M., Meza, J. L., & Wallman, J. (2017). An Analysis of Correlation Between Agricultural Contaminated Wells and Birth Defects in Nebraska.

## Acknowledgement

We want to thank Dr. Tom Rosenquist for assessing morphology of the harvested embryos. This work was funded by an NU Foundation Research Council grant.