



September 2020

## Acute Colonic Perforation in Renal Transplant Recipients: A Case Series

Emily Zurbuchen et al

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>



Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

### Recommended Citation

Zurbuchen, E. Acute Colonic Perforation in Renal Transplant Recipients: A Case Series. Graduate Medical Education Research Journal. 2020 Sep 29; 2(1).  
<https://digitalcommons.unmc.edu/gmerj/vol2/iss1/35>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact [digitalcommons@unmc.edu](mailto:digitalcommons@unmc.edu).

---

## Acute Colonic Perforation in Renal Transplant Recipients: A Case Series

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

**Conclusion:** Contrary to prevailing expectations and our study hypothesis, we found that thiamine supplementation in hospitalized patients with encephalopathy did not improve LOS. Our results are consistent with the lack of beneficial effects of supplemental thiamine in other disease conditions such as sepsis (VITAMINS study) and congestive heart failure.<sup>3-5</sup> We do not recommend routine thiamine supplementation in hospitalized patients. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.033>

## Acute Colonic Perforation in Renal Transplant Recipients: A Case Series

Emily Zurbuchen<sup>1</sup>, Nathalie Sela<sup>2</sup>, Alexander Maskin<sup>1</sup>

<sup>1</sup>University of Nebraska Medical Center, Department of Surgery

<sup>2</sup>University of Nebraska Medical Center, Department of Surgery, Division of Transplantation Surgery

**Mentor:** Alexander Maskin

**Program:** General Surgery

**Type:** Case Report

**Introduction:** There are few cases of non-diverticulitis episodes of colonic perforation following kidney transplantation throughout literature. In the systematic literature review by de'Angelis et al, emergency abdominal surgery following solid organ transplantation for non-diverticulitis causes gastrointestinal perforation accounts of 9.2% of cases, and about 58% of those perforations occurred in the colon. The aim of this case series is to review the events surrounding our two

## References

- 1 Attaluri P, Castillo A, Edriss H, Nugent K. Thiamine Deficiency: An Important Consideration in Critically Ill Patients. *Am J Med Sci.* 2018;356(4):382-390.
- 2 Isenberg-Grzeda E, Kutner HE, Nicolson SE. Wernicke-Korsakoff-syndrome: under-recognized and under-treated. *Psychosomatics.* 2012;53(6):507-16
- 3 Fujii T, Luethi N, Young PJ, et al. Effect of vitamin C, hydrocortisone, and thiamine vs hydrocortisone alone on time alive and free of vasopressor support among patients with septic shock: The VITAMINS randomized clinical trial. *JAMA.* 2020.
- 4 Kattoor AJ, Goel A, Mehta JL. Thiamine therapy for heart failure: A promise or fiction? *Cardiovasc Drugs Ther.* 2018;32(4):313-317.
- 5 Smithline HA, Donnino M, Blank FSJ, et al. Supplemental thiamine for the treatment of acute heart failure syndrome: A randomized controlled trial. *BMC Complement Altern Med.* 2019;19(1):96-019-2506-8.

cases of colonic perforation following renal transplantation in hopes to gain understanding of this rare occurrence. Consent was obtained from the patients.

**Case:** Since 2011, there have been two events of transverse colonic perforation in the acute post-operative period following renal transplantation occurring at this institution. Large volume intraperitoneal air observed on plain film upon work up gastrointestinal symptoms prompted urgent surgical intervention in both cases. Both patients underwent urgent exploratory laparotomy and Hartmann procedure with resection of perforated transverse colon and proximal ostomy. The first patient underwent colostomy

takedown and primary colonic re-anastomosis about one year after his kidney transplant. The second patient is currently scheduled for future colostomy reversal.

**Conclusion:** Various theories of non-diverticular causes of colonic perforations are described in literature including cytomegalovirus enterocolitis, uremic enterocolitis, fecal impaction, and corticosteroid administration among others. Despite the lack of clear cause of perforation, it is imperative to have a high index of suspicion for these immunocompromised patients for prompt surgical treatment. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.034>

Proceedings of the 2nd Annual Graduate Medical Education Research Symposium | Poster Presentations

## Does Prophylactic Administration of Tranexamic Acid Reduce Mean Operative Time and Perioperative Blood Loss in Posterior Cervical Spinal Fusion Surgery?

Emmett J. Gannon<sup>1</sup>, Evan P. Larson<sup>1</sup>, Zachary C. Bailey<sup>1</sup>, Sydney Powers<sup>2</sup>, Scott A. Vincent<sup>1</sup>, Chris A. Cornett<sup>1</sup>, Liz Lyden<sup>3</sup>

<sup>1</sup>University of Nebraska Medical Center, Department of Orthopaedic Surgery and Rehabilitation

<sup>2</sup>University of Nebraska Medical Center, College of Medicine

<sup>3</sup>University of Nebraska Medical Center, College of Public Health, Department of Biostatistics

**Mentor:** Chris Cornett

**Program:** Orthopaedic Surgery and Rehabilitation

**Type:** Original Research

**Background:** Adequate control of bleeding during spine surgery is paramount to the success and efficiency of the operation. Tranexamic acid (TXA) is a systemic antifibrinolytic that is beneficial in reducing blood loss during various orthopaedic procedures. TXA's role in posterior cervical spinal surgery, however, remains poorly described. Therefore we sought to evaluate

whether prophylactic TXA administration prior to posterior cervical fusion reduces perioperative blood loss and mean operative time.

**Methods:** Patients undergoing three to six-level posterior cervical fusions were retrospectively enrolled into two groups. In the study group, 21 patients received TXA prior to surgery, and in the control group, 21 patients did not. Patient's younger than 19 years old, and those undergoing surgery for infection or tumors were excluded. Intraoperative blood loss, operative time, and postoperative drain output were measured in addition to demographic data.

**Results:** No difference existed between groups in regards to demographics. The study group had a lower mean operative time (94 min) vs the control group (110 min), although not statistically significant ( $p=0.06071$ ). Postoperative day 1 drain output was significantly lower in the study group ( $p<0.0239$ ). Intraoperative blood loss, day 2, 3 and total drain output were lower in the study group but not statistically significant. There were two postoperative hematomas in the control group. No postoperative thromboembolic events occurred in either group.