September 2020

Sleep Improvement Program and Academic Performance

Matthew Tingle

University of Nebraska Medical Center

Liz Wellsandt

University of Nebraska Medical Center

Kari Nelson

University of Nebraska Medical Center

RJ Barber

University of Nebraska Medical Center

Beau Konigsberg

University of Nebraska Medical Center

See next page for additional authors

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>

Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Tingle, M., Wellsandt, L., Nelson, K., Barber, R., Konigsberg, B., Lydiatte, M., Ewald, C., , McManigal, M.
Sleep Improvement Program and Academic Performance. Graduate Medical Education Research Journal.
2020 Sep 29; 2(1).
<https://digitalcommons.unmc.edu/gmerj/vol2/iss1/28>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

Sleep Improvement Program and Academic Performance

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Authors

Matthew Tingle, Liz Wellsandt, Kari Nelson, RJ Barber, Beau Konigsberg, Maxwell Lydiatte, Cale Ewald, and Matthew McManigal

Sleep Improvement Program and Academic Performance

Matthew Tingle¹, Liz Wellsandt², Kari Nelson³, RJ Barber¹, Beau Konigsberg¹, Maxwell Lydiatte⁴, Cale Ewald⁴, Matthew McManigal², Matthew Tao¹

¹University of Nebraska Medical Center, Department of Orthopedic Surgery

²University of Nebraska Medical Center, College of Allied Health Professions, Physical Therapy

³University of Nebraska Medical Center, Office of Graduate Medical Education

⁴University of Nebraska Medical Center, College of Medicine

Mentor: Matthew Tao

Program: Orthopedic Surgery

Type: Original Research

Background: The aim of this study is to investigate the impact of improved sleep quality and extension can have on academic performance in medical students. Additional outcome measures include evaluating mood (Profile of Mood States) and sleepiness (Epworth Sleepiness Scale). Our hypothesis is

that improved sleep quality and extension will significantly improve test scores, mood and daytime sleepiness.

Methods: This is a pilot study involving 20 first-year medical students. They will be randomized into either control or intervention groups. Both groups will be monitored for baseline sleep characteristics. The intervention group will then undergo a training program aimed to improve sleep quality and extension. Subjects will be

monitored via actigraphy devices (Actigraph wGT3x-BT) on the dominant hand at all times as well as daily, subjective sleep logs. The primary outcome will be raw test scores at the typical testing intervals based on the curriculum. ESS questionnaire will be administered at the beginning and end of the study, and the POMS will be administered weekly.

Results and Conclusion: Pending ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.027>

The Clinical Relevance of Cement Volume in Percutaneous Vertebral Augmentation

Ravi G. Mirpuri¹, **Michael J. Warden**¹, Virginia Hardie¹, Madhuri Are¹

¹University of Nebraska Medical Center, Department of Anesthesiology

Mentor: Madhuri Are

Program: Anesthesiology

Type: Original Research

Background: Vertebroplasty and Kyphoplasty are two forms of percutaneous vertebral augmentation (PVA), in which polymethylmethacrylate cement is used to stabilize vertebral compression fractures (VCF). This study sought to evaluate the relationship between cement volume and clinical outcomes, including pain reduction, opioid use, and complication rate.

Methods: Retrospective chart review produced 88 patients who received PVA at a tertiary care outpatient pain clinic. Cement

volume, type of PVA, gender, level (thoracic vs lumbar) were collected, as well as clinical outcomes of numeric pain score (NPS) reduction, opioid percent change (OPC), and complications. Both pre-procedure and post-procedure (between 2-4 weeks) data were collected.

Results: Sixty-four patients (72.7%) had statistically significant NPS reduction of $\geq 50\%$ pain ($p < .001$). The mean NPS reduction was 4.45 ($p < .0001$). Also sixty-three out of seventy-two patients who previously used opioids demonstrated some form of opioid reduction which was significant ($p < .0001$). The mean opioid reduction measured across all baseline opioid users was 48%. No statistically significant

differences in NPS or OPC were seen across gender or level type. Linear regression using pre NPS as a covariate showed cement volume did not have a significant effect on NPS change or opioid reduction ($p = 0.173$ and $p = .299$ respectively). Cement leakage occurred as a complication in 10.2% of the patients, although only one complication was symptomatic.

Conclusion: Cement volume does not correlate with clinical outcomes of NPS reduction or OPC. Furthermore, our study reinforced PVA as a beneficial treatment for reducing pain and opioid consumption in patients with painful VCF. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.028>

Diagnostic Quality and Radiation Dose for Pulmonary Embolism Protocol Computed Tomography: ASIR-V vs. Standard Filtered Back Projection

Lei Yu^{1*}, Antonia Gurney^{1*}, Kelly Dea^{1*}, Neil Hansen¹, Valerie Shostrom², Chandrakanth Are³, Anna Zajicek¹

¹University of Nebraska Medical Center, Department of Radiology

²University of Nebraska Medical Center, College of Public Health, Department of Biostatistics

³University of Nebraska Medical Center, Office of Graduate Medical Education

*authors contributed equally to the work

Mentor: Anna Zajicek

Program: Radiology

Type: Review/Meta-analysis

Background: Pulmonary embolism (PE) is the third leading cause of acute cardiovascular disease, contributes to 250,000 American hospitalizations yearly. CT pulmonary angiography (CTPA) is the first-line imaging choice for suspected PE, however radiation

exposure remains a concern. We compared the Dose-Length Product (DLP) in CTPA performed using adaptive statistical iterative reconstruction – V (ASIR-V) versus older filtered back projection (FBP) technique, while also evaluating the ability of ASIR-V to maintain diagnostic accuracy.

Methods: A retrospective review was performed of all CTPA scans completed for suspected PE during a one-year period at our institution. A total of 1095 studies met criteria

for inclusion. Five-hundred and thirty-three (48.7%) of these were performed on a CT scanner utilizing ASIR-V (GE Healthcare), and 562 (51.3%) were performed on an older CT scanner utilizing standard FBP (Lightspeed VCT, GE Healthcare). Dose-length product (DLP) and demographic data were recorded. Original radiology reports from board-certified radiologists for each exam were reviewed. Diagnostic quality was determined by usage of specified