University of Nebraska
Medical Center
Nebraska Medicine

Graduate Medical Education
Research Journal

Volume 2 | Issue 1

Article 41

September 2020

Giant Hemorrhagic Adrenal Pseudocyst in a Postpartum Mother

Lucas Bohannon

University of Nebraska Medical Center

Jacob Meyer

University of Nebraska Medical Center

Sara Despain

University of Nebraska Medical Center

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>

Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Bohannon, L., Meyer, J., , Despain, S. Giant Hemorrhagic Adrenal Pseudocyst in a Postpartum Mother. Graduate Medical Education Research Journal. 2020 Sep 29; 2(1).
<https://digitalcommons.unmc.edu/gmerj/vol2/iss1/41>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

Giant Hemorrhagic Adrenal Pseudocyst in a Postpartum Mother

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Giant Hemorrhagic Adrenal Pseudocyst in a Postpartum Mother

Lucas Bohannon¹, Jacob Meyer¹, Sara Despain¹

¹University of Nebraska Medical Center, Department of Family Medicine

Mentor: Sara Despain

Program: Family Medicine

Type: Case Report

Background: Adrenal cysts occur more frequently in females than males, hemorrhagic adrenal cysts coinciding with parturition are exceedingly rare. These cysts often present with few or vague symptoms but are important to recognize for management as rupture could result in peritoneal seeding of cancerous tissue or even exsanguination.

Case: A 30-year-old on postpartum day one with left sided abdominal pain, a palpable mass, and 2 gram drop in serum Hgb with no typical source of postpartum hemorrhage. Ultrasound demonstrated a large fluid collection, identified as a hemorrhagic adrenal cyst on CT (Figure 1). The patient was transferred to for potential intervention including aspiration vs surgical removal. After observation her hemoglobin improved, the cyst was deemed stable, and she was scheduled for outpatient biopsy. However, it spontaneously ruptured, and the patient underwent aspiration of free serosanguinous abdominal fluid. She recovered without incident; unfortunately, the cyst reformed and ultimately, she underwent adrenalectomy.

Adrenal cysts are a rare incidental finding of 0.06 - 0.18% of individuals, the

Figure 1. Giant hemorrhagic left adrenal cyst displacing left kidney inferiorly on CT scan PPD1.

overwhelming majority of which are female by a ratio of 3:1. Giant hemorrhagic adrenal cysts in pregnancy are astonishingly rare. Identification and proper management are paramount, because parturition represents an unpredictable situation with a potentially predictable course. Derangements of internal organs during or after delivery could result in rupture or hemorrhage. Recognizing this anticipated disturbance in pregnancy would aid in predicted course and management.

Conclusion: Hemorrhagic adrenal cysts present a difficult conundrum when it comes to management, given the lack of guidance

secondary to their rarity. Conservative treatment is possible; however, providers need to continue monitoring of the cyst for complications, including rupture in the setting of labor and hemorrhage as in this postpartum complication. Primary care providers, especially those in resource poor setting should consider transfer to higher level of care. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.040>

Risk of Prostate Cancer in U.S. Veterans With Rheumatoid Arthritis

Kaitlyn Brittan¹, Punyasha Roul¹, Joshua F. Baker^{2,3}, Brian Sauer^{4,5}, Namrata Singh^{6,7}, Harlan Sayles¹, Grant W. Cannon^{4,5}, Ted R. Mikuls¹, Bryant R. England¹

¹University of Nebraska Medical Center, Department of Internal Medicine, Division of Rheumatology

²Corporal Michael J. Crescenz VA

³University of Pennsylvania

⁴VA Salt Lake City

⁵University of Utah

⁶Iowa City VA

⁷University of Iowa

Mentor: Bryant England

Program: Internal Medicine, Division of Rheumatology

Type: Original Research

Background: Patients with rheumatoid arthritis (RA) are at a small, but significantly increased risk of cancer compared with the general population. This risk varies by cancer type, with findings relative to prostate cancer conflicting. Recognizing the male predominance and large number of cases

available in the Veterans Health System, we estimated the risk of prostate cancer in a national population of male US Veterans with RA.

Methods: We identified male patients with RA within the VA Corporate Data Warehouse (1/1/2000-4/1/2018) using an algorithm adapted based on large systematic reviews. Prostate cancer and prostate cancer death were identified from the VA Oncology Raw Domain (ORD) and the National Death Index (NDI). Prostate cancer incidence rates (IR; per 1,000 pt-yrs) and 95% CIs were

calculated from the index. SIR and 95% CI were calculated from the Surveillance, Epidemiology, and End Results Program (SEER).

Results: There were 56,764 RA patients included in analysis. Over 406,013 pt-yrs of follow-up, there were 1,433 incident prostate cancers (IR 3.53; 95% CI 3.35-3.72 per 1,000 pt-yrs). Relative to SEER rates, SIR of prostate cancer in RA was 0.93 (95% CI 0.88-0.98).