



September 2020

The Effect of Weather on Orthopedic Injury Presentation to the Emergency Department

Ryan Houghton et al.

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>



Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Houghton, R. The Effect of Weather on Orthopedic Injury Presentation to the Emergency Department. Graduate Medical Education Research Journal. 2020 Sep 29; 2(1).
<https://digitalcommons.unmc.edu/gmerj/vol2/iss1/56>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

The Effect of Weather on Orthopedic Injury Presentation to the Emergency Department

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

The Incidence, Prevalence and Burden of Gout in the Veterans Health Administration

Lindsay Helget¹, Bryant England¹, Punyasha Roul¹, Harlan Sayles², Alison Petro¹, Kaleb Michaud¹, Ted Mikuls¹

¹University of Nebraska Medical Center, Department of Internal Medicine, Division of Rheumatology

²University of Nebraska Medical Center, College of Public Health, Department of Biostatistics

Mentor: Ted Mikuls and Bryant England

Program: Internal Medicine, Division of Rheumatology

Type: Original Research

Background: To determine the prevalence, incidence, and burden of gout in the Veterans Health Administration (VHA) from 2005-2014.

Methods: We used national VHA data from 1/1999-12/2014 to determine the annual incidence and prevalence of gout in the VHA. Gout burden to the VHA was determined by the proportion of patients with

an encounter related to gout. Rates of urate lowering therapy (ULT) and opiate use were determined annually. Characteristics of those with and without gout were compared using 2014 data.

Results: From 2005 to 2014, gout prevalence in the VHA increased from 4.2% to 5.8% while disease incidence ranged from 5.8 to 7.4 cases per 1000 patient-years. Gout prevalence was highest among men, older patients, and non-Hispanic blacks. During 2014, 4.0% of all inpatient or outpatient encounters and 1.3% of hospitalizations were gout-related. ULT administration remained stable over the ten-year period, with 46%

of gout patients receiving ULT in 2014. In contrast, 16.4% of prevalent gout patients were receiving a weak opioid in 2014, nearly doubling the prescription rate of weak opioids in 2005 while the use of stronger opioids did not change significantly over this period. Patients with gout had greater comorbidity and healthcare utilization than patients without gout.

Conclusion: The burden posed by gout in the VHA is considerable and increased between 2005 and 2014. While the use of ULT has remained stable, the use of opioid therapy has increased among patients with gout. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.052>

A Beer League Degloving

Hannah Hornsby¹, Hudson McGinnis¹, T. Jason Meredith¹

¹University of Nebraska Medical Center, Department of Family Medicine

Program: Family Medicine

Type: Case Report

Introduction: Morel-Lavallée lesions are rare, closed degloving injuries between the skin and underlying fascia that creates a fluid collection within the pre-fascial plane. Common causes are high-energy, blunt force trauma and crush injuries.

Case: A 29-year-old male who presented with left thigh pain, swelling and ecchymosis after attempting a "pop-up" slide into third base during a softball game. His foot caught the base and he flipped forward on his extended

knee. He had immediate swelling and pain but ambulated off the field. In clinic two days later, he reported increased swelling and bruising but denied loss of strength or instability. Exam showed thigh swelling and large ecchymosis throughout the thigh and into the knee and upper shin. He had reduced flexion but preserved extension. MRI demonstrated a large anteromedial thigh fluid collection, consistent with large Morel-Lavallée lesion. He successfully treated conservatively with tight compression and activity restriction. With serial follow-ups and evaluation with sports medicine, the patient continued to improve.

Discussion: Prompt compressive therapy obviates complications of infection, fluid expansion and tissue necrosis, which when present require more invasive interventions. With small, acute lesions, compression alone often suffices for resolution, however, larger lesions can require percutaneous drainage, sclerodosis or debridement. Our case highlights the need to consider Morel Lavallée lesions in expanding soft tissue injuries, even when not associated with high velocities. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.053>

The Effect of Weather on Orthopedic Injury Presentation to the Emergency Department

Ryan Houghton¹, Liz Lyden¹, Aaron Barksdale¹

¹University of Nebraska Medical Center, Department of Emergency Medicine

Mentor: Aaron Barksdale

Program: Emergency Medicine

Type: Original Research

Background: Predicting patient flow and presentation in the emergency department (ED) is difficult. Chief complaints vary and often appear random from an anecdotal perspective. Prior studies attempting to correlate weather conditions with orthopedic injuries have been conflicting. The purpose of this study was to explore associations between weather patterns and orthopedic presentations to the UNMC ED.

Methods: Retrospective chart review was conducted at a tertiary academic hospital with 64,000 annual ED visits. Utilizing electronic medical records, all patients visiting the ED between 9/1/2012 to 4/30/2019 were included. ICD10 codes identified patients diagnosed with fractures of upper (distal forearm through wrist) and lower (proximal femur and hip) extremities. Daily weather patterns were obtained through the National Oceanic and Atmospheric Administration, specifically daily temperature, amount and type of precipitation (none, rain/melted snow, ice/snow). Logistic regression was used to determine if weather conditions were predictive of fractures. Results are displayed as odds ratios and 95%

confidence intervals. P<0.05 was considered statistically significant.

Results: Over an 80-month period 373,409 patients were included, with 4,416 fractures identified. Overall, there was a statistically significant association between snow/ice and fractures (p<0.0001). Specifically, an increased risk of upper extremity fracture (OR 1.47, CI 1.28-1.69) and lower extremity fracture (OR 1.25, CI 1.02-1.53). The risk of upper extremity fracture further increased when snow/ice accumulation was > 3 inches (OR 2.64, CI 1.91-3.65). Rain/melted snow was not associated with increased fractures.

Conclusion: There is a significant association between upper and lower extremity fractures with snow/ice accumulation, but not rain/melted snow. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.054>

The Diagnostic Utility of a Myeloid Mutational Panel for Myelodysplastic Syndromes and Myelodysplastic/Myeloproliferative Neoplasms

Warda Ibrar¹, Jesse Cox¹, Catalina Amador¹, Hina N. Qureishi¹, Kai Fu¹, Timothy C. Greiner¹, Allison Vokoun¹, Weiwei Zhang¹, Yuan Ji¹

¹University of Nebraska Medical Center, Department of Pathology

Mentor: Yuan Ji

Program: Pathology, Division of Hematopathology

Type: Original Research

Background: The diagnosis of myelodysplastic syndromes (MDS) and myelodysplastic/myeloproliferative neoplasms (MDS/MPN) is based on morphology and cytogenetics/FISH findings per 2017 WHO classification. With rare exceptions (e.g. SF3B1 mutations), somatic mutations have not been incorporated as diagnostic criteria. In this study we analyzed the utility of mutational analysis with a targeted 54-gene or 40-gene next generation sequencing (NGS) panel in diagnosis of MDS and MDS/MPN.

Methods: The clinicopathologic data and NGS results of patients with unexplained cytopenia with or without cytosis who underwent a bone marrow (BM) biopsy and had sequencing with either a 54 gene Trusight Myeloid or a 40 gene OncoPrint Myeloid NGS Assay performed at our institution from 2017 to 2019. Morphologic diagnosis of BM biopsy based on 2017 WHO classification was considered gold standard.

Results: A total 74 patients were identified, including 25 low-grade MDS (peripheral

blood blasts <1% and BM blasts <5%), 16 high-grade MDS (peripheral blood blasts 2-19% and BM blasts 5-19%), 5 therapy-related MDS, 14 MDS/MPN, and 14 morphological negative ones. Of 74 patients, 158 somatic mutations involving 37 genes were detected and had variant allele frequency (VAF) ranging from 3% to 99%. 62% (23/37) genes showed recurrent mutations and 38% (14/37) genes had one mutation each. The most common mutated genes were TET2, ASXL1, RUNX1, SF3B1 and TP53. Morphological negative, low-grade MDS, high-grade MDS, therapy-related MDS and MDS/MPN showed an average number of somatic mutations with a mean VAF: 1.5/20%, 2/37%, 3.1/38%, 2/33% and 4.8/41%, respectively. Mutations in TP53, RUNX1, or ASXL1 were associated with high-grade and therapy-related MDS (p=0.004), while SF3B1 mutations were associated with low-grade

MDS (p=0.001). In 60 of 74 patients with a diagnosis of MDS or MDS/MPN, 32 showed abnormal cytogenetics and 28 showed normal cytogenetics. One or more mutations were detected in 25 of 28 (89%) MDS or MDS/MPN patients with normal cytogenetics. The sensitivity, specificity, and positive predictive value (PPV) and negative predictive value (NPV) of mutations for MDS and MDS/MPN with different cutoffs are summarized in Table 1.

Conclusion: A myeloid mutational panel provides additional evidence of clonality besides cytogenetics/FISH studies in diagnosis of cytopenia with or without cytosis, and ≥ 2 mutations with ≥ 10% VAF highly predicts MDS and MDS/MPN with a PPV of 100%. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.055>

Table 1.

Diagnostic performance of mutations for MDS and MDS/MPN with different cutoffs.

	Any mutations (VAF≥1%)	VAF≥20%	≥2 mutations	VAF≥10% and ≥2 mutations	VAF≥10% and ≥2 mutations, or abnormal cytogenetics/FISH
Sensitivity	87%	75%	58%	58%	83%
Specificity	50%	93%	93%	100%	86%
PPV	88%	98%	98%	100%	96%
NPV	47%	46%	34%	36%	54%

Coronary Artery Bypass Grafting Versus Percutaneous Coronary Intervention for Left Main Disease in Chronic Kidney Disease Patients

Aravdeep Jhand¹, Raahat Bansal², Rahul Dhwan¹, J. Dawn Brown³, Thomas Porter¹, James Tcheng⁴, Yiannis Chatzizisis¹, Andrew Goldsweig¹

¹University of Nebraska Medical Center, Department of Internal Medicine, Division of Cardiovascular Disease

²Creighton University, Department of Internal Medicine

³Brown University, Department of Internal Medicine, Division of Cardiovascular Disease

⁴Duke University, Department of Internal Medicine, Division of Cardiovascular Disease

Mentor: Andrew Goldsweig

Program: Internal Medicine, Division of Cardiovascular Disease

Type: Review/Meta-analysis

Background: Limited data exists on the optimal revascularization strategy for patients with left main coronary artery disease

(LMCAD) and chronic kidney disease (CKD). We conducted this updated meta-analysis to compare the outcomes of coronary artery bypass graft (CABG) versus percutaneous coronary intervention (PCI) in this population.

Methods: Multiple electronic databases were searched for studies comparing CABG and PCI in CKD patients undergoing

revascularization of LMCAD (> 50% diameter stenosis). CKD was defined as estimated glomerular filtration rate of < 60 ml/min/1.73 m². Primary outcome of interest was long term major adverse cardiovascular and cerebral events (MACCE) which was a composite of cardiovascular death, myocardial infarction (MI), stroke and repeat revascularization. Other outcomes of interest