September 2020

Improving the Efficiency of Same-Day III Calls

Tracy Farrell

University of Nebraska Medical Center

Teri L. Schuldt

University of Nebraska Medical Center

Mary Wells

University of Nebraska Medical Center

R. Gregory Bociek

University of Nebraska Medical Center

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>

Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Farrell, T., Schuldt, T. L., Wells, M., , Bociek, R. Improving the Efficiency of Same-Day III Calls. Graduate Medical Education Research Journal. 2020 Sep 29; 2(1).
<https://digitalcommons.unmc.edu/gmerj/vol2/iss1/85>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

Improving the Efficiency of Same-Day III Calls

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

References

- 1 Goldstein, N. P., Frey, S. M., Fagnano, M., Okelo, S. O., & Halterman, J. S. (2018). Identifying Which Urban Children With Asthma Benefit Most From Clinician Prompting: Subgroup Analyses From the Prompting Asthma Intervention in Rochester—Uniting Parents and Providers (PAIR-UP) Trial. *Academic Pediatrics*, 18(3), 305-309. doi:10.1016/j.acap.2017.08.015
- 2 Khasha R., Sepheri MM, Mahdavi SA. (2019). An ensemble learning method for asthma control level detection with leveraging medical knowledge-based classifier and supervised learning. *Journal of Medical Systems*, 43(6):158. doi: 10.1007/s10916-019-1259-8
- 3 Rojanasart S and Carlson AM. (2017). The Medical Home Model and Pediatric Asthma Symptom Severity: Evidence from a National Health Survey. *Population Health Management*, 21(2):130-138. doi: 10.1089/pop.2017.0066
- 4 Ruffner MA, Henrickson SE, Chilutti M, Grundmeier R, Spergel JM, Brown-Whitehorn TF. (2018). Improving office scheduling increases patient follow up and reduces asthma readmission after pediatric asthma hospitalization. *American College of Asthma, Allergy, and Immunology*, 121(5):561-567. doi: 10.1016/j.anai.2018.08.015
- 5 Sommer SJ, Queenin L.M., Nethersole S, Greenberg J. (2011). Children's hospital Boston community Asthma initiative: partnerships and outcomes advance policy change. – *Prog Community Health Partnersh*, 5(3):327-35. doi:10.1353/cpr.2011.0044

High-Persensitivity Pneumonitis

Daniel Van Kalsbeek¹, Daniel Hershberger²

¹Department of Internal Medicine, University of Nebraska Medical Center,

²Department of Internal Medicine, Division of Pulmonary and Critical Care Medicine, University of Nebraska Medical Center,

Mentor: Daniel Hershberger

Program: Internal Medicine

Type: Case Report

Introduction: Hypersensitivity pneumonitis (HP) is an immune-mediated lung disease characterized by parenchymal inflammation following inhalation of an inciting antigen.

Case Report: A middle-aged woman was admitted with sub-acute respiratory failure. Physical exam revealed a euvolemic woman with coarse breath sounds who was requiring supplemental oxygen. She had two prior hospitalizations for similar but less severe

symptoms. Her previous episodes appeared to respond to antibiotics though required her to go home on oxygen. Chest CT revealed diffuse bilateral consolidations, and a leukocytosis was present. Broad spectrum antibiotics were initiated. There was no clinical improvement at 24 hours so TMP/SMX plus corticosteroids were added to her treatment plan. She promptly improved. A hypersensitivity panel was positive for antibodies against *Aspergillus* species. Exposure history uncovered marijuana use that coincided with her current and prior respiratory symptoms. Hypersensitivity pneumonitis was diagnosed. She was discharged on a steroid taper and strict marijuana avoidance. At six-week follow

up her symptoms had completely resolved.

Discussion: Environmental fungi are well-established triggers for HP. They are able to pass through lit marijuana cigarettes and pipes, particularly the spores of *A. fumigatus*. Knowing this we felt confident making the diagnosis of HP in our patient because of her positive hypersensitivity panel plus her history of marijuana use coinciding with her symptoms. Her symptoms resolved with corticosteroids and abstinence from marijuana.

Conclusion: Marijuana harbors organic antigens and should be explored as a trigger for hypersensitivity pneumonitis. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.083>

Improving the Efficiency of Same-Day Ill Calls

Tracy Farrell¹, Teri L. Schultdt¹, **Mary Wells¹**, R. Gregory Bociek¹

¹University of Nebraska Medical Center, Department of Internal Medicine, Division of Oncology and Hematology

Mentor: R. Gregory Bociek

Program: Internal Medicine, Division of Oncology and Hematology

Type: Original Research

Background: We sought to use quality improvement (QI) methodology to approach problems leading to burnout and workplace stress in our oncology care team.

Methods: Through a broad survey, we identified coordination of same-day, outpatient visits for acutely ill oncology patients as a source of excessive stress for team members across various disciplines including case managers, infusion clinic nurses, advanced practice providers, and

physicians. We used the Plan-Do-Study-Act (PDSA) framework to identify a feasible, appropriate intervention to reduce case manager time required to coordinate visits with the specific aim of reducing their time by 25%. We also tracked phone calls required to arrange a visit and perceived frustration with the process. Our intervention involved transitioning from paging individual practitioners to using HIPAA-compliant group text to collaboratively coordinate care.

Results: After one PDSA cycle, we found case manager time required to arrange a visit had decreased by 21%, number of calls required by 59% and frustration with the process by 41%.

Conclusion: While we did not meet our specific aim, we feel use of a QI approach led to an easily implemented, effective modification to streamline a previously inefficient, disruptive workflow. The use of QI methodology ensured we understood the baseline process and involved all stakeholders before implementing a change and also ensured we followed data to understand our intervention's impact on team members. Viewing sources of burnout and workplace frustration through the lens of QI may lead to more consistently high-yield interventions than traditional wellness-based, administrative approaches. ■

<https://doi.org/10.32873/unmc.dc.gmerj.2.1.084>