October 2021

Hemorrhagic Bullae in the Setting of a Rare Opportunistic Infection

Taylor K. Thieman

University of Nebraska Medical Center

Ritu N. Swali

University of Nebraska Medical Center

Corey J. Georgesen

University of Nebraska Medical Center

Tell us how you used this information in this [short survey](#).

Follow this and additional works at: <https://digitalcommons.unmc.edu/gmerj>

Part of the [Higher Education Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Thieman, T. K., Swali, R. N., , Georgesen, C. J. Hemorrhagic Bullae in the Setting of a Rare Opportunistic Infection. Graduate Medical Education Research Journal. 2021 Oct 04; 3(1).
<https://digitalcommons.unmc.edu/gmerj/vol3/iss1/46>

This Conference Proceeding is brought to you for free and open access by DigitalCommons@UNMC. It has been accepted for inclusion in Graduate Medical Education Research Journal by an authorized editor of DigitalCommons@UNMC. For more information, please contact digitalcommons@unmc.edu.

Hemorrhagic Bullae in the Setting of a Rare Opportunistic Infection

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Hemorrhagic Bullae in the Setting of a Rare Opportunistic Infection

Taylor K. Thieman¹, **Ritu N. Swali**², Corey J. Georgesen²

¹University of Nebraska Medical Center, College of Medicine

²University of Nebraska Medical Center, College of Medicine, Department of Dermatology

Mentor: Corey J. Georgesen

Program: Dermatology

Type: Case Report

Background: Dermatological presentation of infectious disease can be non-specific, with morphology ranging from erythematous plaques and ulcers to vesicles and bullae. In cases of disseminated infection with cutaneous manifestations, skin biopsy can be a comparatively less invasive and more cost-effective bedside diagnostic technique.

Case: We present a case of a 72-year-old woman hospitalized for anemia and hypoxia, with an incidental finding of grouped vesicles on the hand. She then developed worsening lung cavitations and hemorrhagic bullae during the course of admission. Due to the patient's decompensation, lung biopsy was unobtainable, and instead skin biopsy and tissue culture were obtained. Histopathological examination revealed branching gram-positive bacilli, consistent with *Nocardia otitidiscavarium* (Fig. 1).

Conclusion: This case highlights the utility of early dermatologic consultation and skin biopsies as a safe and efficacious tool for diagnosis of complex medical disease. ■

<https://doi.org/10.32873/unmc.dc.gmerj.3.1.021>

Figure 1. Axial non-contrast CT images of the chest showing a large hiatal hernia, multifocal bronchiectasis with multiple small nodules, and tree-in-bud opacities in the mid and lower lung regions. Imaging findings are varied in MAC pulmonary infections, but prominent bronchiectasis is often described.